

Bayer Environmental Science



Before using this product read the label carefully. / Lees die etiket noukeurig alvorens die produk gebruik word.

## CONFIDOR<sup>®</sup> PINS

Reg. No. L8022 Act No. 36 of 1947  
Namibian Reg. No. N-AR 0767

Reg. Nr. L8022 Wet Nr. 36 van 1947  
Namibiese Reg. Nr. N-AR 0767

CONFIDOR PINS IS A READY-TO-USE, SLOW RELEASE INSECTICIDAL PIN FOR THE CONTROL OF APHIDS, MEALYBUGS AND AUSTRALIAN BUGS ON ORNAMENTAL POTPLANTS.

CONFIDOR PINS IS 'N GEREED-VIRGEBRUIK, STADIG VRYSTELLENDSE INSEKODER BEVATTENDE STAFIE VIR DIE BEHEER VAN PLANTLUISE, WITLUISE EN AUSTRALIESE WOLLUISE OP ORNAMENTELE POTPLANTE.

Ornamental potplant treatment / Ornamentele potplant behandeling

### ACTIVE INGREDIENT / AKTIEWE BESTANDEEL:

Imidacloprid (chloro-nicotinyl)                      25 g/kg                      Imidakloprid (chloro-nikotiniel)

NET CONTENTS / NETTO INHOUD: 10x2 g pins/stafies

### REGISTRATION HOLDER / REGISTRASIEHOUER:

Bayer (Pty.) Ltd. / (Edms.) Bpk.  
Reg. No. / Nr. 1968/011192/07  
P.O. Box / Posbus 143  
ISANDO 1600  
SOUTH AFRICA / SUID-AFRIKA  
TEL: (011) 921 5911

Batch no: / Lotnommer:

Date of manufacture: / Datum vervaardig:

CONFIDOR<sup>®</sup> is a registered trademark of Bayer / is 'n geregistreerde handelsmerk van Bayer.



## **WARNINGS:**

Handle with care.  
Harmful when swallowed.  
Harmful to fish and other animals.  
Store in a cool place and away from food and feedstuffs.  
Keep out of the reach of children, uninformed persons and animals.

## **PRECAUTIONS:**

Avoid contact by mouth.  
Wash with soap and water immediately after use.  
Do not eat, drink or smoke whilst applying or before washing hands and face.  
Do not contaminate dams, drains, ponds, streams and other waterways with the product or used containers.  
Destroy empty container and do not re-use for any other purpose.

## **DIRECTIONS FOR USE: USE ONLY AS DIRECTED.**

**Confidor® Pins** is effective against aphids, mealybugs and Australian bugs on potplants and plants in bags.

Determine the number of pins needed as indicated on the table below. Break the pins in half. Push the halved pins an equal distance from one another, at a distance halfway between the plant and the pot rim / bag rim, until the top of the pin is just below the surface of the soil (see figure). Water the plant. Do not water excessively to the point that water flows from the pot / bag, because this water will contain active ingredient which will lead to less effective control of the listed insects.

Effective control can be expected within 7 to 10 days after treatment.

Repeat the treatment if live insects are noted again, about 6 to 8 weeks after initial treatment.

Note: The pin is slow-release and will not dissolve immediately.

<b>Diameter of pot / bag:</b>	<b>Number of pins:</b>
< 10 cm	½ pin
10-14 cm	1 pin
15-20 cm	2-3 pins
> 20 cm	4 pins

For any product enquiry, contact Bayer Environmental Science - 011 921 5911. For poisoning emergencies only, Griffon Poison Information Centre may also be contacted - 082 446 8946.

**CONFIDOR®** is a registered trademark of Bayer.

© Bayer (Pty.) Ltd. All unauthorized reproduction of this label is forbidden.

www.bayergarden.co.za

## **WAARSKUWINGS:**

Hanteer versigtig.  
Skadelik indien ingesluk word.  
Skadelik vir visse en ander diere.  
Berg in 'n koel plek weg van voer en voedsel.  
Hou buite bereik van kinders, oningeligte persone en diere.

## **VOORSORGMAATREËLS:**

Verhoed kontak deur mond.  
Was met seep en water onmiddellik na gebruik.  
Moet nie eet, drink of rook tydens toediening en voordat hande en gesig gewas is nie.  
Moenie damme, rioolstelsels, afvoerpipe, riviere, dammetjies of ander waterweë met die produk of gebruikte houers besoedel nie.  
Vernietig leë houers en moet vir geen doel hergebruik nie.

## **GEBRUIKSAANWYSINGS:** GEBRUIK SLEGS SOOS AANGEDUI.

**Confidor® Pins** is effektief teen plantluise, wolluise en Australiese wolluise in potplante en plante in sakke.

Bepaal die aantal stafies benodig soos aangedui op die onderstaande tabel. Halveer die stafies. Druk die gehalveerde stafies ewe ver van mekaar af, op 'n afstand die helfte tussen die plantstam en rand van die pot / sak totdat die bokant van die stafie net onder die grond is (sien figuur). Benat die plant, maar moenie die plant oorbenat sodat water uit die pot / sak vloei nie. Water wat uit die pot / sak uitspoel sal die aktiewe bestanddeel bevat wat sal lei tot verminderde effektiwiteit teen genoemde insekte.

Effektiewe beheer kan verwag word binne 7 tot 10 dae na behandeling.

Herhaal die behandeling as lewendige insekte weer waargeneem word, ongeveer 6 tot 8 weke na aanvanklike behandeling.

Nota: Die stafie is stadig-vrystellend en sal nie onmiddellik oplos nie.

<b>Deursnee van pot / sak:</b>	<b>Aantal stafies:</b>
< 10 cm	½ stafie
10-14 cm	1 stafie
15-20 cm	2-3 stafies
> 20 cm	4 stafies

Vir enige produk navrae, skakel Bayer Environmental Science - 011 921 5911. Slegs vir vergiftigingsnoodgevalle kan Griffon Gif Inligtingsentrum ook gekontak word - 082 446 8946.

**CONFIDOR®** is 'n geregistreerde handelsmerk van Bayer.

© Bayer (Edms.) Bpk. Alle ongemagtigde kopiëring van die etiket is verbode.

www.bayergarden.co.za