



ULTRAKILL

Rat and Mouse Pellets

REG. NO. L 9322 Act No. 36 of 1947
Namibian Reg. No. N-AR 1603

REG. NR. L 9322 Wet Nr. 36 of 1947
Namibiese Reg. Nr. N-AR 1603

A HIGHLY ACTIVE ANTICOAGULANT
GRAIN BAIT FOR THE CONTROL OF
THE NORWAY RAT (*Rattus norvegicus*),
ROOF RAT (*Rattus rattus*) AND HOUSE
MOUSE (*Mus musculus*)

'N HOOGS AKTIEWE ANTISTOLMIDDEL GRAAN
LOKAAS VIR DIE BEHEER VAN DIE
NOORWEEGSE ROT (*Rattus norvegicus*),
DAKROT (*Rattus rattus*) EN HUISMUIS (*Mus
musculus*)

ACTIVE INGREDIENTS/AKTIEWE BESTANDELE:

Flocoumafen

0.05g/kg

Flokumafen

NET CONTENTS:

REGISTRATION HOLDER

Starplex 83 CC
P.O. Box 14374
BREDELL 1623
SOUTH AFRICA
+27119794246/7

Distribution by:

Coopers Environmental Science (Pty) Ltd
Reg. No. 2000/02366/70/7
P.O. Box 14374
BREDELL 1623
SOUTH AFRICA
TEL: +2711 979-4246/7

Emergency number:

Poison Information Centres: South Africa

Gauteng: Unitas Hospital: 0800 111 9900 (all hours)

Western Cape: Tygerberg Hospital: (021) 931 6129



Harmful
Skadelik



WARNINGS:

Handle with care
Keep out reach of children, animals and uninformed people.
Store away from food and feedstuffs.
Poisonous if swallowed and upon prolonged contact with the skin.
Toxic to fish and wildlife
In case of poisoning call a doctor and make this label available to him.

PRECAUTIONS:

Avoid contact with skin. Wear gloves when handling product.
Avoid contamination of any food, food utensils and drinking water.
Set bait where it will be inaccessible to children and animals.
Wash hands and exposed skin after handling.
Destroy empty container and do not reuse for any other purpose.
Bury dead rodents and any bait not consumed at end of treatment.

SYMPTOMS OF POISONING

Bleeding from the gums and nose, bleeding under the skin, pallor, haemorrhage into and around elbow and knee joints. Blood in urine and faeces.

NOTE TO PHYSICIAN

Administer 5mg/kg Vit. K₁ intravenously on the first day, followed by another administration of the same dose if the quick time does not normalize a few hours after the first injection. Follow up with 5 mg/kg/day orally on the second and following days. The duration of the treatment is generally between 10 and 15 days.

In severe cases give a small transfusion of fresh blood

DIRECTIONS FOR USE: USE ONLY AS DIRECTED.

For use in the home, on the farm, and industrial premises.

ULTRAKILL Rat and Mouse Pellets can kill rats and mice after a single feed. Death of the rodent will occur within 4 -12 days.

Effective against other anticoagulant resistant rats and mice.

Norway and Roof Rats

Place **ULTRAKILL Rat and Mouse Pellets** in a covered bait station such as the **Coopers Coral Bait Station** or a similar model to prevent access by children and domestic animals. Place 50-75g **ULTRAKILL Rat and Mouse Pellets** per bait station every 5 – 10 m along walls or rat trails, in rat burrows and sheltered areas where rats are known to live and feed.

Bait points should be inspected and replenished every 7 days until feeding ceases.

Mice

Place 15-20g **ULTRAKILL Rat and Mouse Pellets** per bait station (such as the **Coopers Peti Bait Station**) every 2 – 4 m along the walls or runways, in nests and in sheltered areas where mice are known to live or feed.

The single feed action of **ULTRAKILL Rat and Mouse Pellets** is particularly effective against mice with their small appetites and sporadic feeding habits.