



MAXFORCE® QUANTUM

Reg. No. L8460 Act No. 36 of 1947
Namibian Reg. No. N-AR 0970

Reg. Nr. L8460 Wet Nr. 36 van 1947
Namibiese Reg. Nr. N-AR 0970

A READY-TO-USE, ODOURLESS, INCONSPICIOUS AND NON-REPELLANT INSECTICIDAL BAIT FOR THE CONTROL OF ANTS IN AND AROUND DOMESTIC PREMISES, PUBLIC BUILDINGS, FOOD STORAGE AND HANDLING PREMISES AND INDUSTRIAL PREMISES.

'N GEREED-VIR-GEbruik, REUKLOSE, ONOPVALLENDE EN NIE-AFWERENDE INSEKODODENDE LOKAAS VIR DIE BEHEER VAN MIERE IN EN RONDOM HUISHOUDELIKE PERSELE, PUBLIEKE GEBOUE, VOEDSEL-OPBERGING EN -HANTERINGSPERSELE, EN INDUSTRIËLE PERSELE.

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL:

Imidacloprid (chloro-nicotinyl)

0.3 g/kg

Imidakloprid (chloro-nikotiniel)

NET CONTENTS: / NETTO INHOUD: 30 g

REGISTRATION HOLDER / REGISTRASIEHOUER:

Bayer (Pty.) Ltd. / (Edms.) Bpk.
Reg. No. / Nr. 1968/011192/07
P.O. Box / Posbus 143
ISANDO 1600
SOUTH AFRICA / SUID-AFRIKA
TEL: (011) 921 5911

Batch no: / Lotnommer:

Date of manufacture: / Datum vervaardig:

MAXFORCE® is a registered trademark of the Bayer / is 'n geregistreerde handelsmerk van die Bayer.



MAXFORCE® QUANTUM

Reg. No. L8460 Act No. 36 of 1947
Namibian Reg. No. N-AR 0970

A ready-to-use, odourless, inconspicuous and non-repellant insecticidal bait for the control of ANTS in and around domestic premises, public buildings, food storage and handling premises and industrial premises.

ACTIVE INGREDIENT: Imidacloprid (chloro-nicotinyl) 0.3 g/kg

REGISTRATION HOLDER:

Bayer (Pty.) Ltd.
Reg. No. 1968/011192/07
P.O. Box 143
ISANDO 1600
SOUTH AFRICA
TEL: (011) 921 5911

WARNINGS:

Handle with care.
Store in a cool place and away from food and feedstuffs.
Keep out of the reach of children, uninformed persons and animals.

PRECAUTIONS:

Avoid contact with skin.
Wear suitable gloves during application.
Wash with soap and water immediately after use or accidental skin contact.
Do not eat, drink or smoke whilst applying or before washing hands and face.
Apply only in positions where it will be completely inaccessible to children and animals.
Do not apply bait to food or feedstuffs or surfaces on which food or feed is stored, prepared or eaten.
Keep bait in original packaging in a cool, safe and dry place. Do not freeze.
Destroy used container and do not re-use for any other purpose.

DIRECTIONS FOR USE: Use only as directed.

Suitable for use in domestic, commercial and industrial premises; food handling, processing and storage establishments, e.g. restaurants, food retailers, warehouses, raw material stores, commercial kitchens and abattoirs; public buildings e.g. hospitals, hotels, municipal buildings, churches, halls, community centres and cinemas; outdoor treatments e.g. terraces, pavements, patios, entrance areas to sheds.

Maxforce® Quantum must be dispensed by using a suitable gel applicator. After application, remove the empty cartridge and dispose of safely. If the cartridge is only partially used, remove it from the applicator and seal it with the end cap provided.

Apply out of reach and sight of children and animals. Place bait near or on ant trails. Survey the area to be treated to identify sites of ant infestation. An effect on the ant population can be expected within 10 days when the frequency of foragers will decline. If not consumed, baiting points will remain effective for up to 12 weeks. Avoid re-invasion by providing baiting points continuously.

INDOOR:

Apply **Maxforce® Quantum** directly as spots in identified target areas. Instruct residents to avoid cleaning or removing bait spots. Where infestation is high, inspect the spots regularly and replenish bait if necessary. Avoid application to excessively dusty, damp or greasy locations and areas which are subject to washing. Do not apply to food preparation surfaces or areas that have recently been treated with other insecticides. Do not contaminate the bait with other chemicals.

Apply 1 spot of 200 mg bait/m of ant trail

OUTDOOR:

Inject bait directly into nest entrances. Repeat when necessary.

Apply 2 g bait per nest entrance

SA Poison Centre: 021 931 6129 / 0800 333 444

MAXFORCE® is a registered trademark of Bayer.

© Bayer (Pty.) Ltd. All unauthorized reproduction of this label is forbidden.

www.bayeres.co.za

MAXFORCE® QUANTUM

Reg. Nr. L8460 Wet Nr. 36 van 1947
Namibiese Reg. Nr. N-AR 0970

'N GEREED-VIR-GEBRUIK, REUKLOSE, ONOPVALLENDE EN NIE-AFWERENDE INSEKODODENDE LOKAAS VIR DIE BEHEER VAN MIERE IN EN RONDOM HUISHOUDELIKE PERSELE, PUBLIEKE GEBOUE, VOEDSEL-OPBERGING EN -HANTERINGSPERSELE EN INDUSTRIËLE PERSELE.

AKTIEWE BESTANDDEEL: Imidaklopried (chloro-nikotiniel) 0.3 g/kg

REGISTRASIEHOUER:

BAYER (EDMS.) BPK.
Reg. Nr. 1968/011192/07
Posbus 143
ISANDO 1600
SUID-AFRIKA
TEL: (011) 921 5911

WAARSKUWINGS:

Hanteer versigtig.
Berg in 'n koel plek weg van voedsel en voer.
Hou buite bereik van kinders, oningeligte persone en diere.

VOORSORGMATREËLS:

Verhoed kontak met vel.
Dra geskikte handskoene tydens aanwending.
Was met seep en water onmiddellik na gebruik en toevallige velkontak.
Moet nie eet, drink of rook tydens toediening of voordat hande en gesig gewas is nie.
Wend slegs aan op plekke wat totaal onbereikbaar is vir kinders en diere.
Moenie lokaas aanwend op voedsel of voer of oppervlakke waar voedsel of voer geberg, voorberei of geëet word nie.
Berg lokaas in oorspronklike verpakking in 'n koel, veilige en droë plek. Moenie vries nie.
Vernietig leë houers. Moet vir geen doel hergebruik nie.

GEBRUIKSAANWYSINGS: Gebruik slegs soos aangedui.

Geskik vir gebruik in huishoudelike-, handel- en nywerheidspersone; voedselverwerkings-, vervaardigings- en voedselstoorpersone, bv. restaurante, kleinhandelaars, pakstore, rou materiaalstore, kommersiële kombuise en slagpale; publieke geboue bv. hospitale, hotelle, munisipale geboue, kerke, sale, gemeenskapsentrums en bioskope; buitenshuise behandeling bv. terrasse, plaveisel, patio's, ingange na skure, ens.

Maxforce® Quantum moet toegedien word met 'n geskikte jel toediener. Verwyder die leë spuit na toediening en vernietig op 'n veilige wyse. As die spuit nog produk bevat, verwyder dit van die toediener en seël dit met die voorsiene doppie.

Dien toe buite bereik van kinders en diere. Plaas lokaas naby of op mierpaadjies. Ondersoek die area wat behandel moet word om areas van mierbesmetting te bepaal. 'n Effek op die mierbevolking kan verwag word binne 10 dae wanneer die frekwensie van werkers sal afneem. Lokaaspunte sal effektief bly vir 12 weke indien dit nie geneem is nie. Verhoed herbesmettings deur lokaaspunte deurlopend te voorsien.

BINNENSHUIS:

Dien **Maxforce® Quantum** direk toe as kolbehandeling in die geïdentifiseerde teikenareas. Vra inwoners om nie die area skoon te maak of die lokaasdruppels te verwyder nie. Op plekke waar

besmetting hoog is, ondersoek die area gereeld en vul die lokaas aan indien nodig. Vermoed toediening op oormatig stowwerige, klam of vetterige areas wat gereeld gewas word nie. Moenie toedien op voedselvoorbereidingsoppervlakke of areas wat onlangs behandel is met ander insekdoders nie. Moenie die lokaas kontamineer met ander chemikalieë nie.

Dien 1 kol van 200 mg lokaas/m mierpaadjie toe

BUITENSHUIS:

Sput lokaas direk in nesopeninge in. Herhaal wanneer nodig.

Dien 2 g lokaas per nesingang toe

SA Gifsentrum: 021 931 6129 / 0800 333 444

MAXFORCE® is geregistreerde handelsmerk van Bayer.

© Bayer (Edms.) Pty. Alle onregmatige kopieëring van die etiket is verbode.

www.bayeres.co.za