

WARNINGS:

Handle with care.

Poisonous when swallowed or inhaled or absorbed through the skin.

Toxic to bees, fish and wildlife.

Keep out of reach of children, uninformed persons and animals.

Flammable - do not store or use near open flame.

Store away from food and feedstuffs. Do not use on alkaline surfaces or where staining is undesirable.

IN CASE OF POISONING CALL A DOCTOR AND MAKE THIS LABEL AVAILABLE TO HIM.

PRECAUTIONS:

Do not inhale spray mist.

Wash with soap and water immediately after accidental skin contact.

Do not eat, drink or smoke whilst applying and mixing, before washing hands and face.

Wear a face mask and protective clothing when applying insecticide for long periods.

Prevent contamination of food, feedstuffs, drinking water and eating utensils.

Rinse empty container three times with a volume of water equal to at least one tenth of that of the container and add the rinsing's to the contents of the spray tank before disposing of the container.

Destroy empty container and do not re-use for any other purpose.

SYMPTOMS OF POISONING:

Headache, fatigue, faintness, giddiness, excessive sweating, nausea, abdominal pains, vomiting, diarrhea, tightness of the chest, anxiety, blurred vision, muscle twitching beginning in the eyelids and tongue, small pupils, respiratory distress, convulsions, coma.

FIRST AID TREATMENT:

Remove patient from source of poisoning to a well-ventilated area and keep him quiet and reassured.

Remove contaminated clothing and rinse contaminated body area thoroughly with plenty of soap and water.

Do not rub skin. Flush contamination out of eyes with clean water for 15 minutes. If water diluted mixture has been swallowed induce vomiting by tickling the back part of the throat. If concentrate has been swallowed firstly give copious quantities of either beaten egg white, starch solution or Milk of Magnesia before inducing vomiting. Repeat until vomit fluid is clear and free from smell of poison. Administer artificial respiration or closed chest cardiac massage if necessary. Do not apply direct mouth to mouth respiration. Never give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN:

Administer atropine sulphate, intravenously (1 - 4 mg) every 5 - 10 minutes until signs of atropinization (dry flushed skin and tachycardia) appear. 2 PAM 50 mg/kg up to total dose of 1 - 2 g or 2/3 of this dose of Protopam. Repeat oximes if necessary.

DO NOT GIVE MORPHINE.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED.

RESKOL[®] 64 EC must be diluted with water immediately before use.

RESKOL[®] 64 EC gives residual protection.

1. Shake container thoroughly.

2. Cockroaches

Dilute 500 ml/ 10 l water. Apply as a coarse surface spray. Repeat when necessary.

3. Bedbugs

- a) For knockdown add 30 ml/ 10 l water. Repeat when necessary.
- b) For longer protection, add 60 ml/ 10 l water. Repeat when necessary.
- c) Apply as a coarse surface spray, or brush the mixture onto the wall surface at the rate of 5 l/ 100 m², particularly into cracks and crevices. Spray or brush bed frames, under tables, cupboards (not food cupboards), etc.

4. Pests of skins and hides (Dermestidae, e.g. *Dermestes maculatus*)

For protection against these pests use RESKOL[®] 64 EC at the rate of 30 ml/ 10 l water. For longer protection use at the rate of 60 ml/ 10 l water.

Ten liters of prepared wash will treat approximately 100 skins or 24 hides.

5. Flies and mosquitoes

Add 30 ml/ 10 l water. Apply as a coarse surface spray at the rate of 5 l/ 100 m².

Repeat when necessary

WAARSKUWINGS:

Hanteer versigtig.

Giftig wanneer ingesluk, ingesem of deur die vel geabsorbeer word.

Giftig vir visse, bye en natuurlewe.

Hou buite bereik van kinders, oningeligte persone en diere.

Vlambaar - moenie naby oop vlamme gebruik nie.

Bewaar weg van voedsel en voer. Moet nie op alkaliese oppervlaktes of waar verkleuring van die oppervlakte ongewens is, aanwend nie.

IN DIE GEVAL VAN VERGIFTIGING ONTBIED 'N GENEESHEER EN STEL HIERDIE ETIKET AAN HOM BESKIKBAAR.

VOORSORGMAATREËLS:

Moenie sproeiwel inasem nie.

Was onmiddellik met seep en water indien dit per abuis op die vel gekom het.

Moenie eet, drink of rook tydens aanwending of vermenging, alvorens hande en gesig gewas is nie.

Dra 'n gesigsmasker en beskermende klere indien insektemiddel vir lang periodes aangewend word.

Voorkom besoedeling van voedsel, voer, drinkwater en eetgerei.

Spoel leë houer drie keer uit met 'n volume water gelykstaande aan ten minste een tiende van die inhoud van die houer. Gooi die uitspoelwater by die inhoud van die spuitkan voordat die houer vernietig word. Vernietig leë houer en moet vir geen ander doel gebruik nie.

SIMPTOME VAN VERGIFTIGING:

Hoofpyn, vermoeidheid, kragteloosheid, duiseligheid, oormatige sweet, mislikheid, maagpyn, braking, buikloop, benoude bors, angstigheid, versteurde gesigsvermoë, spiertrekkings wat by die ooglede en tong begin, verkleinde pupille, asemhalingsnood, stuiptrekkings, bewusteloosheid.

NOODHULP BEHANDELING:

Neem pasiënt weg van bron van vergiftiging na 'n goed geventileerde omgewing en hou die pasiënt stil en gerusgestel. Verwyder besoedelde klere en spoel die besmette liggaamsdele deeglik af met baie koue water en seep. Moenie die vel vryf nie. Spoel besmette oë met skoon water uit vir 15 minute. Indien van die waterverdunde mengsel ingesluk is, veroorsaak braking deur agter in die keel met vinger te prikkel. As die konsentraat gedrink is, gee eerstens oorvloedige hoeveelhede geklitste eierwit, styseloplossing of Melk van Magnesia, alvorens braking teweeggebring word. Hou hiermee aan totdat die braaksel skoon is en nie meer na die gif ruik nie. Pas kunsmatige asemhaling of geslote hartmassering toe indien nodig. Moenie direkte mond tot mond asemhaling toepas nie. Moet nooit enigiets deur die mond aan 'n bewustelose persoon toedien nie.

NOTA AAN GENEESHEER:

Dien atropien sulfaat binne-aars toe (1 - 4 mg) elke 5 - 10 minute totdat tekens van atropinisasie intree (droë gloeiende vel en tachycardea). 2 PAM 50 mg/kg tot 'n totale dosis van 1 - 2 g of 2/3 van hierdie dosis Protopam. Herhaal toediening van oksieme indien nodig.

MOET GEEN MORFIEN TOEDIEN NIE.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI.

RESKOL[®] 64 EC moet met water verdun word onmiddellik voor gebruik.

RESKOL[®] 64 EC **gee nawerkende beskerming.**

1. Skud die houer deeglik.

2. Kakkerlakke

Voeg 500 ml/ 10 l water. Dien toe as 'n growwe oppervlakbespuiting. Herhaal wanneer nodig.

3 Weeluisse

- a) Om hulle uit te klop voeg 30 ml/ 10 l water. Herhaal wanneer nodig.
- b) Vir langer nawerking voeg 60 ml/ 10 l water. Herhaal wanneer nodig.
- c) Dien toe as growwe bespuiting aan oppervlaktes of wend aan met 'n kwas op die muur teen 5 l/ 100 m², veral in die krakies en barste. Bespuit of wend met 'n kwas aan op slaapbankrame, onder tafels, kaste (nie koskaste nie), ens.

4. Plae van velle en huide (Dermestidae, bv. *Dermestes maculatus*)

Vir beskerming teen hierdie plae moet RESKOL[®] 64 EC gebruik word teen 'n dosis 30 ml/ 10 l water.

Om langer beskerming te verkry moet 60 ml/ 10 l water gebruik word.

Tien liter van die aangemaakte vloeistof is genoeg om ongeveer 100 velle of 24 huide te behandel.

5. Vlieë en muskiete

Voeg 30 ml/ 10 l water. Dien toe as 'n growwe oppervlakbespuiting teen 5 l/ 100 m².

Herhaal wanneer nodig

Vir enige produk navrae, skakel Coopers Environmental Science (Pty) Ltd - 011 979 4246/7. Slegs vir vergiftigingsnoodgevalle kan Griffon Gif Inligtingsentrum ook gekontak word - 082 446 8946.