

Insecticide

NuFog Dual Action Fogger

A total release fogger treatment for the control of crawling and flying insects with the additional benefit of limited residual activity. For use in homes, kitchens, restaurants, hotels and industrial settings.

- ▶ Quick knock-down
- ▶ Total release fogger
- ▶ Good residual action
- ▶ Short exposure time
- ▶ Range of insects controlled



 **COOPERS**
environmental science

www.cooperses.co.za

ACTIVE INGREDIENT:

Cyfluthrin (pyrethroid) 0,25 g/kg
Propoxur (carbamate) 10,0 g/kg
Dichlorvos (organophosphate) 5,0 g/kg

Harmful

For use in:

- Domestic
- Industrial

CONTROLS: Ants, bedbugs, carpet beetles, cockroaches, crickets, fleas, fish moths, house flies and mosquitoes.

PACKAGING: 325 ml

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

APPLICATION: 160m³ per 325ml can

WARNINGS: May cause irritation of the skin, eyes, mucous membranes and skin sensitization. Keep out of reach of children, animals and uninformed persons. Contents under pressure. Flammable – do not spray near open flame or onto hot surfaces. Do not expose to heat exceeding 50 oC or leave in direct sunlight. Do not pierce or incinerate can, even when empty. Ensure good ventilation before re-entry after recommended treatment period.

PRECAUTIONS: Do not use in the presence of persons with known sensitivity to pyrethroids. In the case of signs of sensitivity, obtain medical advice. Prevent direct contamination of food, feedstuff, drinking water and utensils. Avoid excessive inhalation and skin contact. Remove pets and seal aquaria airtight during treatment. May stain plastic material and highly polished surfaces. r any other purpose but destroy it by perforation and flattening and dispose of it in a safe way. Prevent contamination of food, feedstuffs, drinking water and eating utensils.

FIRST AID TREATMENT

Remove patient from source of poisoning to a well-ventilated area and keep him quiet and reassured. Remove contaminated clothing and rinse contaminated body area thoroughly with plenty of soap and water. Flush contamination out of eyes with clean water and call a physician. If ingested, take the patient to the nearest physician immediately.

NOTE TO PHYSICIAN: Treat symptomatically.


DO NOT DEPRESS ACTUATOR UNTIL ALL CANS HAVE BEEN PLACED AND YOU ARE READY FOR COMPLETE TREATMENT ACTIVATION.

Seal off all ventilation to the treatment area. Close all doors and windows of the area to be treated, except the exit door, which should be closed after exiting the area. All applicable precautionary measures should be taken prior to treatment activation. Ensure that all cupboards, cabinets and doors are opened to allow infiltration of the product.

Recommended dosage: One 325 ml can per 160 m³

IMPORTANT: Once the actuator is pressed into the locked position, total content will be released.

1. Place can on newspaper, in the center of the recommended treatment area. Cover surfaces close to the NuFog can with layers of newspaper, or refuse bags.
2. To activate: Point nozzle away from yourself and depress actuator until it locks in position. Start from the furthest room working your way backwards to the exit. Set fogger upright in the center of the newspaper. Immediately leave the treatment area, closing the last door behind you. Keep treated area closed for at least two hours.
3. Ventilate the area well after the treatment period by opening doors and windows, before re-occupation.



 **COOPERS**
environmental science

www.cooperses.co.za