



ProRoach Gel

REG. NO. L 10115 Act No. 36 of 1947

REG. NR. L 10115 Wet Nr. 36 van 1947

INSECTICIDAL GEL BAIT FOR THE CONTROL OF GERMAN AND AMERICAN COCKROACHES (NYMPHS AND ADULTS) IN PUBLIC HYGIENE, DOMESTIC PREMISES AND FOOD STORAGE PREMISES, WHICH IS, NON-REPELLENT ODOURLESS AND NON-VOLATILE.

INSEKDODENDE JEL LOKAAS VIR DIE BEHEER VAN DUITSE EN AMERIKAANSE KAKKERLAKKE (NIMFE EN VOLWASSENES) IN PUBLIEKE HYGIËNE, VOEDSELOPBERGING PERSELE EN HUISHOUDELIKE PERSELE WAT, NIE-AFWEREND, REUKLOOS EN NIE-VLUGTIG IS.

ACTIVE INGREDIENTS/AKTIEWE BESTANDDELE:

Fipronil

0.5 g/kg

Fipronil

NET CONTENTS: / NETTO INHOUD: Contact supplier. 35g

REGISTRATION HOLDER

Starplex 83 CC
P.O. Box 14374
BREDELL 1623
SOUTH AFRICA

Distribution by:

Coopers Environmental Science (Pty) Ltd
Reg. No. 2000/02366/70/7
P.O. Box 14374
BREDELL 1623
SOUTH AFRICA
TEL: +2711 979-4246/7

Emergency numbers:

Poison Information Centres: South Africa
Gauteng: Unitas Hospital: 0800 111 9900 (all hours)
Western Cape: Tygerberg Hospital: (021) 931 6129
For product enquiries, contact Coopers Environmental Science (Pty) Ltd – 086 030 3003. For poisoning emergencies only, contact the Griffon Poison Information Centre - 082 446 8946.



WARNINGS:

Handle with care.

Contact should be avoided by skin and mouth. Harmful if swallowed.

Keep container away from children, uninformed persons and animals.

Store where it is cool, dry and where food and feedstuffs are absent.

PRECAUTIONS:

Exposed skin and hands should be washed after completion and before a meal.

Apply only in positions inaccessible to children or pets.

Do not eat, drink or smoke during application.

Wear gloves during application.

The gel should not be applied onto surfaces where foodstuffs are eaten, stored or prepared.

Contamination of food preparing equipment and utensils must be prevented.

When empty dispose of syringe safely. Do not re-use the syringe for any other purpose.

Waterways, ditches or ponds must not be contaminated with chemical or used container.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED.

General directions

PROROACH GEL an effective gel bait against nymphs and adults of the German cockroach (*Blattella germanica*) and the American cockroach (*Periplanata americana*).

PROROACH GEL may be used to control cockroaches in domestic (kitchens, animal housing), commercial and industrial (food manufacturing premises, commercial kitchens, restaurants, food stores, warehouses, factories, retail outlets, ships etc.) as well as public buildings such as hotels, hospitals, prisons, schools, churches, theatres etc.

Application

For maximum effect, **PROROACH GEL** should be placed within or close to cockroach harbourages, foraging and feeding areas. These are typically areas that are dark, warm, damp and remain undisturbed e.g. behind refrigerators and in electrical equipment. These areas should be surveyed to identify centers of cockroach infestations e.g. by using sticky traps such as the **Ultra Roach Trap** and/or flush-out aerosols such as **PCO Flushing Agent**; searching for droppings, cast skins and egg cases.

For this product to achieve its maximum potential, make sure the premises is thoroughly cleaned before application, removing all other food sources. Surfaces should be thoroughly dried before application.

PROROACH GEL tubes are designed to be used with models such as the **Eco-Gel** or **Airofog Gel Applicators**.

PROROACH GEL should be applied throughout the infested area in many small spots. Spots of gel should be applied in dark out-of-sight locations in voids, cracks, and crevices, corners lift shafts, service-ducts, furniture and equipment e.g. refrigerators, counters, sinks, cookers and around sensitive situations e.g. electrical, mechanical and electronic equipment. **PROROACH GEL** should not be applied **in** areas where it will come into contact with water or be removed by routine cleaning. Dusty, damp or greasy locations should also be avoided. The spraying of other insecticides on or around **PROROACH GEL** spots is not recommended as also the placement of spots on recently treated surfaces, as this will contaminate the **PROROACH GEL** and may discourage cockroaches from feeding on it.

TOEDIENINGDHOEVEELHEDE:

Normal rate	High rate
German cockroach 1 spot of 0,03 g gel /m ² .	German cockroach 2 spots of 0,03 g gel /m ² .
American cockroach 2 spots of 0,03 g gel /m ² .	American cockroach 3 spots of 0,03 g gel /m ² .
A spot of 0,03 g is approximately 3 - 4 mm in diameter.	
Critical Comments: Floor space should be determined to calculate the correct amount of spots that should be placed in the treatment area. Spots should not be placed necessarily uniform, but rather concentrated in areas of high activity such as harbourages. Thus, if a building is 100 m ² and the cockroach is identified as the German cockroach, then 100 spots should be placed around areas of high activity at the normal rate.	

For any product enquiry, contact Coopers Environmental Science (Pty) Ltd - 011 979 4246/7. For poisoning emergencies only, Griffon Poison Information Centre may also be contacted - 082 446 8946.

WAARSKUWINGS:

Stoor weg van kinders, oningesigte persone en diere.
Kontak moet vermy word deur die vel en mond.
Vel irritasie kan voorkom.
Vries van produk moet vermy .
Skadelik vir visse.
Hanteer versigtig.
Suur waar dit koel, droog en waar voedsel en voer afwesig is.

VOORSORGMAATREËLS:

Blootgestelde vel en hande moet met seep en warm water gewas word na voltooiing en voor 'n maaltyd.
Moet nie eet, drink of rook tydens aanwending nie.
Dra handskoene tydens toediening.
Wend slegs aan in toepaslike en veilige areas wat ontoeganklik is vir kinders of troeteldiere.
Moet nie jol aanwend op oppervlaktes waar voer geberg of voedsel geeet of voorberei word nie.
Kontaminasie van voedselvoorbereiding toerusting en gereedskap moet voorkom word.
Na gebruik, raak veilig van lee spuit ontslae. Moet nie spuit vir enige ander doel gebruik nie.
Waterweë, slotte of poele moet nie met chemikaliee of gebruikte houer besoedel word nie.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

ALGEMENE AANWYSINGS:

PROROACH Gel 'n effektiewe gel aas teen nimfe en volwassenes van die Duitse kakkerlak (*Blatella germanica*) en die Amerikaanse kakkerlak (*Periplanata americana*).

PROROACH GEL mag gebruik word kakkerlakte in die binnelandse (kombuise, dier behuising), kommersiële en industriële (voedselverwerkingsaanlegte, kommersiële kombuise, restaurante, winkels, pakhuise, fabrieke, winkels, skepe ens) sowel as openbare geboue soos te beheer hotelle, hospitale, gevangenis, skole, kerke, teaters ens

AANWENDING:

Vir maksimum effek, moet PROROACH GEL binne geplaas word of naby die kakkerlak skuilings, kos soek en voed gebiede. Dit is tipies gebiede wat donker, warm, klam en bly ongestoord by agter yskaste en in elektriese toerusting. Hierdie gebiede moet ondervra word sentrums van kakkerlak besmetting by te identifiseer deur die gebruik van taai lokvalle soos die Ultra Roach Trap en / of spoel-out sputibussoos PCO Flushing Agent; soek vir mis, cast velle en eier gevalle.

Vir hierdie produk sy maksimum potensiaal te bereik, maak seker dat die perseel deeglik skoongemaak word voor toediening, die verwydering van alle ander voedsel bronre. Oppervlaktes moet deeglik droog word voor toediening.

PROROACH GEL buise is ontwerp om gebruik te word met modelle soos die Eco-Gel of Airofog Gel Toedieners.

PROROACH GEL moet dwarsdeur die besmette area toegepas word in baie klein kolle. Plekke van gel moet in die donker buite-oë plekke in ruimtes, krake, en barste toegepas word, hoeke hyserskagte, diens-kanale, meubels en toerusting by yskaste, tellers, wasbakke, kook en om sensitiewe situasies by elektriese, meganiese en elektroniese toerusting. PROROACH GEL moet nie in gebiede waar dit in aanraking kom met water of deur roetine skoonmaak verwijder toegepas word. Stowweringe, klam of vetterige plekke moet ook vermy word. Die bespuiting van ander insekdoders op of rondom PROROACH GEL kolle word nie aanbeveel asook die plasing van die kolle op het onlangs behandel oppervlaktes, want dit sal die PROROACH GEL besoedel en mag van voeding op dit ontmoedig kakkerlakte.

Normale dosis	Hoë dosis
Duitse kakkerlak 1 kol van 0,03 g jel /m ² .	Duitse kakkerlak 2 kolle of 0,03 g jel /m ² .
Amerikaanse kakkerlak 2 kolle van 0,03 g jel /m ² .	Amerikaanse kakkerlak 3 kolle of 0,03 g jel /m ² .
'n Kol van 0,03 g is ongeveer 3 - 4 mm in deursnee.	
Kritiese Kommentaar: Vloeroppervlak moet bepaal word om die korrekte aantal kolle wat in die behandelingsarea geplaas moet word, te bereken. Kolle moet nie noodwendig uniform geplaas word nie, maar eerder gekonsentreer in gebiede van hoë aktiwiteit soos wegkruipplekke. Dus, as 'n gebou 100 m ² groot is en die kakkerlak geïdentifiseer is as die Duitse kakkerlak, dan moet 100 kolle rondom die gebiede van hoë aktiwiteit geplaas word teen die normale koers.	

Vir enige produk navrae, skakel Coopers Environmental Science (Pty) Ltd - 011 979 4246/7. Slegs vir vergiftigingsnoodgevalle kan Griffon Gif Inligtingsentrum ook gekontak word - 082 446 8946.