



# ULTRAKILL Rat and Mouse Grain Bait

REG. NO. L 9188 Act No. 36 of 1947  
Namibian Reg. No. N-AR 1601

REG. NR. L 9188 Wet Nr. 36 of 1947  
Namibiese Reg. Nr. N-AR 1601

**READ THE LABEL BEFORE USE  
KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

A HIGHLY ACTIVE ANTICOAGULANT  
GRAIN BAIT FOR THE CONTROL OF  
THE NORWAY RAT (*Rattus norvegicus*),  
ROOF RAT (*Rattus rattus*) AND HOUSE  
MOUSE (*Mus musculus*)

'N HOOGS AKTIEWE ANTISTOLMIDDEL GRAAN  
LOKAAS VIR DIE BEHEER VAN DIE  
NOORWEEGSE ROT (*Rattus norvegicus*),  
DAKROT (*Rattus rattus*) EN HUISMUIS (*Mus  
musculus*)



**WARNING**

**Hazard Statement:**

- May cause damage to organs (blood) through prolonged or repeated exposure.

**Precautionary Statements:**

- Wash hands and face thoroughly after handling.
- Do not eat, drink or smoke when using this product.
- Dispose of content/container to suitable landfill in accordance with local regulations.

**ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL:**

Flocoumafen 0.05 g/kg

NET CONTENTS:

**REGISTRATION HOLDER**

Starplex 83 CC  
P.O. Box 14374  
BREDELL 1623  
SOUTH AFRICA  
+27119794246/7

**Distribution by:**

Coopers Environmental Science (Pty) Ltd  
Reg. No. 2000/02366/70/7  
P.O. Box 14374  
BREDELL 1623  
SOUTH AFRICA  
TEL: +2711 979-4246/7

**IN CASE OF POISONING CALL THE FOLLOWING NUMBER: (+27) 82 446 8946**

**UN NUMBER: NOT REGULATED**

**WARNINGS:**

Handle with care.  
Keep out reach of children, uninformed people and animals.  
Store away from food and feedstuffs.  
May cause damage to organs (blood) through prolonged or repeated exposure.  
Toxic to wildlife.  
In case of poisoning call a doctor and make this label available to him.

### **PRECAUTIONS:**

Avoid contact with skin. Wear gloves when handling product.  
Do not eat, drink or smoke when using this product.  
Avoid contamination of any food, food utensils and drinking water.  
Set bait where it will be inaccessible to children and animals.  
Wash hands, face and exposed skin after handling.  
Dispose of content/container to suitable landfill in accordance with local regulations. Do not reuse for any other purpose.  
Bury dead rodents and any bait not consumed at end of treatment.

### **RELEVANT SUBSTANCES:**

All relevant hazardous substances within the formulation:  
Flocoumafen 0.005%  
Denatonium benzoate < 1%

### **FIRST AID:**

If inhaled: Keep patient calm, remove to fresh air and seek medical attention.  
Skin contact: Wash thoroughly with soap and water.  
Eye contact: Immediately wash affected eyes for at least 15 minutes under running water with eyelids held open. Consult an eye specialist if necessary.  
If ingested: Rinse mouth immediately and then drink plenty of water and seek medical attention. Never induce vomiting or give anything by mouth if the victim is unconscious or having convulsions. Do not induce vomiting unless told to by a poison control center or doctor.

### **SYMPTOMS OF POISONING**

Bleeding from the gums and nose, bleeding under the skin, pallor, haemorrhage into and around elbow and knee joints. Blood in urine and faeces.

### **NOTE TO PHYSICIAN**

Administer 5 mg/kg Vit. K<sub>1</sub> intravenously on the first day, followed by another administration of the same dose if the quick time does not normalize a few hours after the first injection. Follow up with 5 mg/kg/day orally on the second and following days. The duration of the treatment is generally between 10 and 15 days.

In severe cases give a small transfusion of fresh blood

### **DIRECTIONS FOR USE: USE ONLY AS DIRECTED.**

For use in the home, on the farm, and industrial premises.

**ULTRAKILL Rat and Mouse Grain Bait** can kill rats and mice after a single feed. Death of the rodent will occur within 4 -12 days.

Effective against other anticoagulant resistant rats and mice.

### **Norway and Roof Rats**

Place **ULTRAKILL Rat and Mouse Grain Bait** in a covered bait station such as the **Coopers Charlie Bait Station** or a similar model to prevent access by children and domestic animals. Place 50 – 75 g **ULTRAKILL Rat and Mouse Grain Bait** per bait station every 5 – 10 m along walls or rat trails, in rat burrows and sheltered areas where rats are known to live and feed.

Bait points should be inspected and replenished every 7 days until feeding ceases.

#### Mice

Place 15 – 20 g **ULTRAKILL Rat and Mouse Grain Bait** per bait station (such as the **Coopers Jerry Bait Station**) every 2 – 4 m along the walls or runways, in nests and in sheltered areas where mice are known to live or feed.

The single feed action of **ULTRAKILL Rat and Mouse Grain Bait** is particularly effective against mice with their small appetites and sporadic feeding habits.