



## ProRoach Gel

REG. NO. L 10115 Act No. 36 of 1947

REG. NR. L 10115 Wet Nr. 36 van 1947

**READ THE LABEL BEFORE USE  
KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

GROUP

2B

INSECTICIDE

INSECTICIDAL GEL BAIT FOR THE CONTROL OF GERMAN AND AMERICAN COCKROACHES (NYMPHS AND ADULTS) IN PUBLIC HYGIENE, DOMESTIC PREMISES AND FOOD STORAGE PREMISES, WHICH IS, NON-REPELLENT ODOURLESS AND NON-VOLATILE.

INSEKODENDENDE JEL LOKAAS VIR DIE BEHEER VAN DUITSE EN AMERIKAANSE KAKKERLAKKE (NIMFE EN VOLWASSENES) IN PUBLIEKE HYGIËNE, VOEDSELOPBERGING PERSELE EN HUISHOUDELIKE PERSELE WAT, NIE-AFWEREND, REUKLOOS EN NIE-VLUGTIG IS.



**DANGER**

#### Hazard Statements:

- Causes damage to organs through prolonged or repeated exposure.
- Very toxic to aquatic life with long lasting effects.

#### Precautionary Statements:

- Do not breathe mist, vapours and spray.
- Wash hands and face thoroughly after handling.
- Do not eat, drink or smoke when using this product.
- Avoid release into the environment.
- Collect spillage.
- Dispose of content/container to suitable landfill in accordance with local regulations.

#### ACTIVE INGREDIENTS/AKTIEWE BESTANDDELE:

Fipronil

0.5 g/kg

Fipronil

NET CONTENTS: / NETTO INHOUD: Contact supplier. 35g

#### REGISTRATION HOLDER

Starplex 83 CC  
P.O. Box 14374  
BREDELL 1623  
SOUTH AFRICA

**IN CASE OF POISONING CALL THE FOLLOWING NUMBER: (+27) 82 446 8946**

**UN NUMBER: 3077**

## **WARNINGS:**

Handle with care.  
Contact should be avoided by skin and mouth.  
Causes damage to organs through prolonged or repeated exposure.  
Store where it is cool, dry and where food and feedstuffs are absent.  
Very toxic to aquatic life with long lasting effects.

## **PRECAUTIONS:**

Wash hands and face thoroughly after handling.  
Apply only in positions inaccessible to children or pets.  
Do not eat, drink or smoke when using this product.  
Wear gloves during application.  
The gel should not be applied onto surfaces where foodstuffs are eaten, stored or prepared.  
Contamination of food preparing equipment and utensils must be prevented.  
When empty dispose of syringe safely. Do not re-use the syringe for any other purpose.  
Collect spillage. Avoid release to the environment. Waterways, ditches or ponds must not be contaminated with chemical or used container.  
Dispose of content/container to suitable landfill in accordance with local regulations.

## **RELEVANT SUBSTANCES:**

All relevant hazardous substances within the formulation:

0.05% Fipronil  
< 1% Methylparaben  
< 10% Isopropylalcohol

## **FIRST AID:**

IF INHALED: Keep patient calm, remove to fresh air, and seek medical attention.  
ON SKIN CONTACT: After contact with skin, wash immediately with plenty of water and soap. If irritation develops, seek medical attention.  
ON CONTACT WITH EYES: Immediately wash affected eyes for at least 15 minutes under running water with eyelids held open, consult an eye specialist.  
IF INGESTED: Immediately rinse mouth and then drink plenty of water, do not induce vomiting, seek medical attention.

**DIRECTIONS FOR USE:** USE ONLY AS DIRECTED.

### **General directions**

**PROROACH GEL** an effective gel bait, containing a stomach poison, for control against nymphs and adults of the German cockroach (*Blattella germanica*) and the American cockroach (*Periplaneta americana*).

**PROROACH GEL** is suitable for use in domestic areas (kitchens, animal housing), commercial and industrial (food manufacturing premises, commercial kitchens, restaurants, food stores, warehouses, factories, retail outlets, ships etc.) as well as public buildings such as hotels, hospitals, prisons, schools, churches, theatres etc.

### **Application**

For **PROROACH GEL** to achieve its maximum potential, make sure the premises is thoroughly cleaned before application, removing all other food sources. Surfaces should be dry before application.

For maximum effect, **PROROACH GEL** should be placed within or as close to cockroach harbourages, foraging and feeding areas as possible. These areas should be surveyed beforehand to identify infestation hotspots (search for droppings, cast-off skins, and discarded egg cases.) The use of sticky traps such as the **Ultra Roach Trap** and/or flush-out aerosols such as **PCO Flushing Agent** is advised.

**PROROACH GEL** cartridges are designed to be used with the **Eco-Gel**, **Airofog Gel** or similar gel applicators.

**PROROACH GEL** is most effective when many small bait points are applied throughout the infested area, concentrating on hotspots such as harbourages (see below table). It is still necessary to determine the number of bait points needed by using the dosage table below, these bait points can then be concentrated on the identified areas.

In general, gel bait points should be applied in dark concealed locations such as in cracks, and crevices, corners lift shafts, service-ducts, voids, furniture and equipment e.g. refrigerators, counters, sinks, cookers and around sensitive situations e.g. electrical, mechanical and electronic equipment. Moist areas or areas that are often cleaned should be avoided as bait placement areas. Dusty, damp or greasy locations should also be avoided. Do not spray other insecticides on or around **PROROACH GEL** bait points, as this will contaminate the **PROROACH GEL** and could minimize consumption by cockroach.

**APPLICATION RATES:**

<b>Moderate infestation</b>	<b>High infestation</b>
German cockroach <b>1 bait point</b> of 0,03 g gel /m <sup>2</sup> .	German cockroach <b>2 bait points</b> of 0,03 g gel /m <sup>2</sup> .
American cockroach <b>2 bait points</b> of 0,03 g gel /m <sup>2</sup> .	American cockroach <b>3 bait points</b> of 0,03 g gel /m <sup>2</sup> .
A bait point of 0,03 g is approximately 3 - 4 mm in diameter.	